

# Spatio-temporal and ontogenetic changes in the diet of the Argentine hake *Merluccius hubbsi*

MAURO BELLEGGIA<sup>1,2,3</sup>, DANIEL E. FIGUEROA<sup>2</sup>, GABRIELA IRUSTA<sup>1</sup> AND CLAUDIA BREMEC<sup>1,3</sup>

<sup>1</sup>Instituto Nacional de Investigación y Desarrollo Pesquero, Paseo Victoria Ocampo 1, Mar del Plata, B7602HSA, Argentina,

<sup>2</sup>Laboratorio de Ictiología, Instituto de Investigaciones Marinas y Costeras, Universidad Nacional de Mar del Plata, Funes 3350, B7602AYL, Mar del Plata, Argentina, <sup>3</sup>Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Avenida Rivadavia 1917, C1033AAJ, Ciudad Autónoma de Buenos Aires, Argentina

*The diet and the trophic level of the Argentine hake Merluccius hubbsi was investigated in the south-west Atlantic Ocean. Stomachs from 3405 specimens, ranging from 10 to 94 cm total length, collected on three research cruises carried out during winter 2011 and summer 2012, were examined at the laboratory. Dietary shifts related to sex, maturity stage, size, age, season, region and hour were examined using generalized linear models. The Argentine hake M. hubbsi fed mainly on zooplankton crustaceans (Themisto gaudichaudii, Euphausia lucens and Munida spp.), followed by fish and cephalopods. During summer, cephalopods (mainly Illex argentinus) were the main prey of M. hubbsi, indicating spatio-temporal changes in the diet. This work also revealed ontogenetic dietary changes associated with size, from zooplankton crustaceans during early life stages (E. lucens, T. gaudichaudii and Munida spp.) to fish and cephalopods. The trophic level of M. hubbsi ranged from 3.43 to 4.51 according to season and region. Cannibalism increased with predator length and contributed a maximum of 8.14% to the diet of M. hubbsi during winter on the inner Patagonian continental shelf.*

**Keywords:** feeding ecology, trophic level, Patagonia, Merlucciidae

Submitted 24 January 2014; accepted 4 April 2014

## INTRODUCTION

The family Merlucciidae comprises three subfamilies: Merlucciinae, Steindachneriinae and Macruroninae (Nelson, 2006). The former includes one genus, *Merluccius*, with 13 recognized species (Nelson, 2006). The common name applied to these species is hake (Nelson, 2006), and they usually have commercial value throughout the world (Lloris *et al.*, 2003). The Argentine hake *Merluccius hubbsi* is endemic to the south-west Atlantic and has a latitudinal range from Brazil (21°30'S) to southern Patagonia (55°S) (Cousseau & Perrotta, 2004). It is the most important ground-fish resource found in the Argentine sea (Bezzi *et al.*, 1995; Pitcher & Alheit, 1995).

There are two main stocks of *M. hubbsi* on the Argentinean continental shelf (ACS), the northern stock (NS, 34–41°S) and the Patagonian stock (PS, 41–55°S) (Bezzi *et al.*, 1995). A third small stock is located in the semi-enclosed San Matías Gulf (Di Giacomo *et al.*, 1993). These stocks have different historical exploitation patterns and are managed as separated units (Aubone *et al.*, 2004). The PS exhibits a well known migration pattern that could be summarized as winter offshore and summer inshore (Macchi *et al.*, 2007). During early summer, hake move from deeper waters (more

than 110 m depth) to the coast (close to the 50 m isobath) for spawning (Macchi *et al.*, 2007). Spawning begins in November–December at 50 m depth at ~43–44°S. After spawning, the hake return to deeper waters where they remain broadly dispersed throughout the open sea during winter on the Patagonian continental shelf (Macchi *et al.*, 2007).

The Argentine hake is an abundant species and an important predator on the Patagonian shelf waters (Laptikhovsky & Fetisov, 1999). Although numerous studies have focused on the diet of the Argentine hake on the ACS in the past decades (Angelescu *et al.*, 1958; Angelescu & Cousseau, 1969; Cordo, 1981; Angelescu & Prenschi, 1987; Sánchez & Prenschi 1996; Ruiz & Fondacaro, 1997), on different stocks (Sánchez & García de la Rosa, 1999; Sánchez, 2009; Ocampo Reinaldo *et al.*, 2011) or on certain age-classes (Temperoni *et al.*, 2013), feeding studies conducted at large spatio-temporal scales including many variables are lacking. For instance, the diet of *M. hubbsi* on the Malvinas Islands shelf still remains unknown (Arkhipkin *et al.*, 2003). Moreover, it is necessary to update the trophic role of the Argentine hake, especially in the PS where the community has been subjected to high fishing pressure during the last decades (Aubone *et al.*, 2004). The aims of this study were to: (1) provide a quantitative taxonomic description of the diet of the Argentine hake on the PS; (2) analyse the influence of sex, maturity stage, size, age, season, region and hour on their diet; and (3) determine the trophic level of the species.

**Corresponding author:**

M. Belleggia

Email: belleggia@inidep.edu.ar

## MATERIALS AND METHODS

### Study area and sample collection

The study area covered the Patagonian continental shelf, between 41°S and 48°S, and from 57 m to 144 m depth (Figure 1). The Patagonian continental shelf is washed offshore by the Malvinas Current, which extends northward from Drake Passage (~55°S) to nearly 38°S (Bastida *et al.*, 1992; Piola *et al.*, 2010). This subantarctic water enters the shelf, mainly between the Malvinas Islands and Tierra del Fuego, and is diluted by continental discharges (Guerrero & Piola, 1997). The mean current flow on the Patagonian continental shelf is from south to north (Guerrero & Piola, 1997). A tidal mixing front, located at 43–45°S during spring and summer, defines the boundary between stratified (offshore) waters and a coastal, vertically mixed body of water (Acha *et al.*, 2004). The spawning of Argentine hake in the Patagonian stock is primarily associated with this thermal front (Ehrlich & Ciechowski, 1994).

The Argentine hake of the PS exhibits seasonal migrations, moving into inshore areas in summer at 43–45°S for spawning (Macchi *et al.*, 2007). The specimens disperse after spawning, moving back to deeper waters during the cold season (Macchi *et al.*, 2007). Based on this spatio-temporal *a priori* knowledge, the study area was subdivided into the inner Patagonian continental shelf (IPCS; depth <110 m, latitude >43°S, longitude >61°W) and the outer Patagonian continental shelf (OPCS; depth >110 m or latitude <43°S and longitude >61°W) (Figure 1). During winter, the Argentine hake is homogeneously distributed along both the IPCS and the OPCS (Macchi *et al.*, 2007). During summer, the species aggregates for spawning on the IPCS (Macchi *et al.*, 2007). Given the known migration pattern and the spawning aggregations during summer, the variables season and region were combined and categorized into the following spatio-temporal




Fig. 1. Map of the study area showing the sample sites where specimens of *Merluccius hubbsi* were collected during winter on inner Patagonian continental shelf (winter inner Patagonian continental shelf (IPCS), black circles), winter on outer Patagonian continental shelf (winter outer Patagonian continental shelf, crosses) and summer on inner Patagonian continental shelf (summer IPCS, white circles).

variable (SPTP): winter IPCS, winter OPCS and summer IPCS (Figure 1).

Specimens were caught during two trawling surveys conducted by the Instituto Nacional de Investigación y Desarrollo Pesquero (INIDEP, Argentina) during July, August and September 2011 and January 2012. Specimens caught between July and early September were regarded as having been caught during austral winter and those caught in January represented those caught in austral summer. Fishing was conducted during daylight (07.00–19.00 hours), at 3–4 knots for 30 min at each sampling site, using an Engel type bottom trawl net (200 mm mesh in the wings, 103 mm in the cod ends, 4 m vertical opening and 15 m horizontal aperture). Specimens captured were measured to the nearest cm and sexed. Sexual maturity (mature or immature) was determined for females according to the condition of ovary and the presence of yolk eggs, whereas male maturity was determined by the size and colour of testes (Macchi & Pájaro, 2003). The sagittae otoliths were collected for age determination following the method proposed by Renzi & Pérez (1992). Finally, stomachs were excised, labelled in plastic bags and frozen at –25°C for analysis at the laboratory.

### Diet composition and data analysis

In the laboratory, prey items found in the stomachs were identified to the lowest possible taxonomic level using taxonomic works (Ramírez, 1971; Bastida & Torti, 1973; Ramirez & Viñas, 1985), field guides (Cousseau & Perrota, 2004) and by consulting specialists. Prey items were counted and weighed to the nearest 0.01 g. The contribution of each prey to the diet was evaluated by calculating the percentage frequency of occurrence (%F, the total number of stomachs in which a given prey was found expressed as percentage of the total number of stomachs with food), the numerical percentage (%N, the total number of a given prey as percentage of the total number of prey found) and the percentage of weight (%W, the weight of a given prey as percentage of the total weight of prey found). These three parameters were integrated into the index of relative importance:  $IRI = \%F \times (\%N + \%W)$  (Pinkas *et al.*, 1971), expressed as a percentage (%IRI; Cortés, 1997).

The trophic level (TrL) was estimated for the species following the method proposed by Cortés (1999) as follows:

$$TrL = 1 + \left( \sum_{j=1}^n P_j \times TrL_j \right)$$

where  $TrL_j$  is the trophic level of each prey item  $j$  and  $P_j$  is the proportion of each prey item  $j$  (using %IRI) in the diet of *M. hubbsi*, and  $n$  is the total number of prey items. The  $TrL_j$  were obtained from the literature (Table 1; Ebert & Bizzarro, 2007; Vögler *et al.*, 2009; Sea Around Us, 2014).

### Dietary models

The numbers of the six main prey items of the Argentine hake (*Themisto gaudichaudii*, *Euphausia lucens*, fish, cephalopods, *Munida* spp. (includes *M. gregaria* and *M. subrugosa*) and *Merluccius hubbsi* (cannibalism)) were used as dependent variables. This kind of numerical data has many zeros, has a variance much greater than the mean and exhibits a negative

**Table 1.** Diet composition of the Argentine hake *Merluccius hubbsi*, on the winter inner Patagonian continental shelf (winter inner Patagonian continental shelf (IPCS)), winter outer Patagonian continental shelf (winter outer Patagonian continental shelf (OPCS)) and summer inner Patagonian continental shelf (summer IPCS). The diet is expressed by the percentage frequency of occurrence (%F), the percentage of number (%N), percentage of weight (%W) and the percentage of index of relative importance (%IRI). TrL<sub>j</sub> is the trophic level of each prey obtained from: Ebert & Bizzarro, 2007; Vögler *et al.*, 2009; and Sea Around Us, 2014.

Prey items	Winter IPCS				Winter OPCS				Summer IPCS				TrL <sub>j</sub>
	%F	%N	%W	%IRI	%F	%N	%W	%IRI	%F	%N	%W	%IRI	
<b>Fish</b>	<b>20.863</b>	<b>1.291</b>	<b>68.590</b>	<b>12.081</b>	<b>29.684</b>	<b>0.922</b>	<b>56.822</b>	<b>15.059</b>	<b>15.673</b>	<b>1.670</b>	<b>7.349</b>	<b>1.615</b>	
Merlucciidae <i>Merluccius hubbsi</i>	8.39	0.37	56.06	8.14	4.42	0.08	12.80	0.86	3.39	0.33	2.16	0.16	4.08 b)
Merlucciidae <i>Macruronus magellanicus</i>	–	–	–	–	0.21	<0.01	3.80	0.01	–	–	–	–	3.9 c)
Nototheniidae <i>Patagonotothen ramsayi</i>	8.99	0.73	10.59	1.75	14.32	0.32	18.16	3.97	2.46	0.31	0.51	0.04	3.49 b)
Clupeidae <i>Sprattus fuegensis</i>	0.84	0.07	0.33	0.01	–	–	–	–	–	–	–	–	3.4 c)
Engraulidae <i>Engraulis anchoita</i>	0.36	0.02	1.06	0.01	5.89	0.14	14.87	1.33	8.77	0.93	4.35	0.88	2.48 b)
Zoarcidae <i>Ilucoetes fimbriatus</i>	–	–	–	–	–	–	–	–	0.12	0.01	0.16	<0.01	3.24 a)
Ophidiidae <i>Genypterus blacodes</i>	0.12	0.01	0.04	<0.01	0.21	<0.01	6.08	0.02	–	–	–	–	4.34 b)
Ophidiidae <i>Raneya brasiliensis</i>	–	–	–	–	–	–	–	–	0.12	0.01	0.11	<0.01	3.56 b)
Myxinidae	0.12	0.01	0.21	<0.01	–	–	–	–	–	–	–	–	3.24 a)
Non-identified fish	2.16	0.09	0.31	0.01	5.89	0.10	1.10	0.11	0.82	0.08	0.05	<0.01	3.24 a)
<b>Crustaceans</b>	<b>87.41</b>	<b>98.413</b>	<b>22.428</b>	<b>87.528</b>	<b>77.263</b>	<b>98.984</b>	<b>22.852</b>	<b>82.703</b>	<b>44.561</b>	<b>91.716</b>	<b>4.457</b>	<b>48.959</b>	
Euphausiidae <i>Euphausia lucens</i>	41.61	57.19	5.43	44.76	53.89	86.44	14.95	82.04	21.17	24.95	0.19	10.16	2.25 a)
Hyperiididae <i>Themisto gaudichaudii</i>	58.03	38.30	5.77	43.95	42.32	11.56	2.72	9.07	12.16	27.75	0.12	6.47	3.18 a)
Galatheidae <i>Munida gregaria</i>	4.80	0.92	4.60	0.45	1.68	0.61	3.10	0.09	7.37	26.08	2.47	4.02	2.52 a)
Galatheidae <i>Munida subrugosa</i>	1.56	0.18	2.34	0.07	0.42	0.01	0.03	<0.01	0.58	0.06	0.04	<0.01	2.52 a)
Galatheidae <i>Munida</i> spp.	7.07	0.79	2.26	0.37	8.84	0.37	2.04	0.32	8.89	11.16	1.31	2.11	2.52 a)
Sergestidae <i>Peisos petrunkevitchi</i>	6.24	0.73	1.08	0.19	–	–	–	–	3.74	1.52	0.18	0.12	2.4 a)
Solenoceridae <i>Pleoticus muelleri</i>	1.80	0.15	0.47	0.02	–	–	–	–	1.05	0.10	0.13	<0.01	2.52 a)
Pandalidae <i>Austropandalus grayi</i>	1.44	0.09	0.07	<0.01	–	–	–	–	–	–	–	–	2.52 a)
Squillidae <i>Pterygosquilla armata</i>	0.24	0.01	0.40	<0.01	–	–	–	–	0.23	0.03	0.03	<0.01	2.52 a)
Majidae <i>Libidoclaea granaria</i>	0.12	0.01	0.01	<0.01	–	–	–	–	–	–	–	–	2.52 a)
Gammaridae	0.36	0.02	<0.01	<0.01	0.21	0.01	<0.01	<0.01	0.23	0.05	<0.01	<0.01	3.18 a)
Isopoda <i>Acanthoserolis</i> spp.	0.12	0.01	<0.01	<0.01	–	–	–	–	–	–	–	–	3.18 a)
Isopoda <i>Cirolana</i> spp.	0.12	0.01	<0.01	<0.01	–	–	–	–	–	–	–	–	3.18 a)
Non-identified crustaceans	0.24	0.02	<0.01	<0.01	–	–	–	–	0.23	0.02	<0.01	<0.01	2.52 a)
<b>Cephalopods</b>	<b>5.156</b>	<b>0.281</b>	<b>8.864</b>	<b>0.391</b>	<b>12.421</b>	<b>0.349</b>	<b>20.152</b>	<b>2.241</b>	<b>45.731</b>	<b>6.533</b>	<b>88.070</b>	<b>49.424</b>	
Ommastrephidae <i>Illex argentinus</i>	0.12	0.01	4.90	0.01	1.68	0.03	7.61	0.19	43.04	6.25	86.15	75.93	3.8 c)
Loliginidae <i>Loligo</i> spp.	3.84	0.22	3.46	0.24	10.32	0.31	12.49	1.99	2.11	0.24	1.89	0.09	3.2 a)
Sepiolidae <i>Semirossia</i> spp.	1.20	0.06	0.51	0.01	0.42	0.01	0.05	<0.01	0.23	0.02	<0.01	<0.01	3.2 a)
Non-identified cephalopods	–	–	–	–	–	–	–	–	0.23	0.02	0.02	<0.01	3.2 a)
<b>Others</b>	<b>0.36</b>	<b>0.015</b>	<b>0.117</b>	<b>&lt;0.01</b>	<b>0.632</b>	<b>0.010</b>	<b>0.174</b>	<b>0.001</b>	<b>0.819</b>	<b>0.080</b>	<b>0.125</b>	<b>0.002</b>	
Polychaeta <i>Chaetopterus</i> spp.	–	–	–	–	–	–	–	–	0.12	0.01	0.05	<0.01	2.6 a)
Polychaeta <i>Phyllochaetopterus</i> spp.	0.12	0.01	0.06	<0.01	–	–	–	–	–	–	–	–	2.6 a)
Polychaeta tube	0.12	0.01	0.03	<0.01	–	–	–	–	0.35	0.03	0.01	<0.01	2.6 a)
Non-identified polychaeta	–	–	–	–	–	–	–	–	0.12	0.01	<0.01	<0.01	2.6 a)
Porifera <i>Tedania</i> spp.	0.12	0.01	0.02	<0.01	0.63	0.01	0.17	<0.01	0.12	0.01	<0.01	<0.01	2.5 a)
Echinodermata <i>Cosmasterias lurida</i>	–	–	–	–	–	–	–	–	0.12	0.01	0.07	<0.01	2.5 a)

binomial distribution (Crawley, 2005). All these issues are dealt with by using the generalized linear model (GLM) with negative binomial error distribution and a log link (Crawley, 2005). The independent variables selected to explain the consumption of the main prey were: sex; maturity stage (juvenile, adults); total length (TL, cm); age; SPTP (winter IPCS, winter OPCS and summer IPCS); and hour of the day (07.00–19.00). In order to test a curvilinear relationship between TL and the consumption of any prey, models including a quadratic term of TL ( $TL^2$ ) as independent were also fitted (Table 2). The models with all possible combinations of two independent variables were constructed (Table 2). A theoretical model without an independent variable was also constructed to test the hypothesis that none of the independent variables selected in this work influenced the consumption of any prey (Table 2) (Lucifora *et al.*, 2009). Akaike's information criterion (AIC) and Akaike's weight ( $w$ ) were used to compare and rank a total of the 28 competing models, and to estimate which of them best fit the data (Franklin *et al.*, 2001; Johnson & Omland, 2004). The model with the lowest AIC was selected and plotted (Franklin *et al.*, 2001; Johnson & Omland, 2004). When a quadratic relationship was selected for the consumption of any prey, the trend was evaluated by fitting LOESS regressions (Crawley, 2005). The LOESS fits a local quadratic regression model using non-parametric techniques to produce a smoothed model surface (Crawley, 2005). All analyses were performed using R program (<http://www.R-project.org>).

**Table 2.** List of the generalized linear models (GLM) fitted to explain the consumption of the main prey of the Argentine hake *Merluccius hubbsi*. The independent variables selected are: sex; maturity, maturity stage; TL, total length; TL +  $TL^2$ , quadratic total length; age; SPTP, spatio-temporal; hour, hour of the day; and 1, null model.


Models					
GLM 1	Prey	~	Sex		
GLM 2	Prey	~	Maturity		
GLM 3	Prey	~	TL		
GLM 4	Prey	~	Age		
GLM 5	Prey	~	SPTP		
GLM 6	Prey	~	Hour		
GLM 7	Prey	~	TL + $TL^2$		
GLM 8	Prey	~	Sex	+	Maturity
GLM 9	Prey	~	Sex	+	TL
GLM 10	Prey	~	Sex	+	Age
GLM 11	Prey	~	Sex	+	SPTP
GLM 12	Prey	~	Sex	+	Hour
GLM 13	Prey	~	Sex	+	TL + $TL^2$
GLM 14	Prey	~	Maturity	+	TL
GLM 15	Prey	~	Maturity	+	Age
GLM 16	Prey	~	Maturity	+	SPTP
GLM 17	Prey	~	Maturity	+	Hour
GLM 18	Prey	~	Maturity	+	TL + $TL^2$
GLM 19	Prey	~	TL	+	Age
GLM 20	Prey	~	TL	+	SPTP
GLM 21	Prey	~	TL	+	Hour
GLM 22	Prey	~	Age	+	SPTP
GLM 23	Prey	~	Age	+	Hour
GLM 24	Prey	~	Age	+	TL + $TL^2$
GLM 25	Prey	~	SPTP	+	Hour
GLM 26	Prey	~	SPTP	+	TL + $TL^2$
GLM 27	Prey	~	Hour	+	TL + $TL^2$
GLM 28	Prey	~	1		

## RESULTS


### Diet composition

During winter in the IPCS, 1187 specimens ranging from 10 cm to 83 cm TL were examined (Figure 2), of which 834 (70.26%) contained prey items in their stomachs. *Merluccius hubbsi* fed mainly on crustaceans (Figure 3; Table 1). The euphausiid *Euphausia lucens* was the first in importance by %IRI and %N, and the second by %F (Table 1). The hyperiid amphipod *Themisto gaudichaudii* was the most important prey item in terms of %F, and the second one by %IRI and %N (Table 1). The gadoid *M. hubbsi* (by cannibalism) and the notothenioid *Patagonotothen ramsayi* were the most important prey in terms of %W (Table 1). The trophic level (TrL) of the Argentine hake *M. hubbsi* during winter in the IPCS was 3.84.

During winter in the OPCS, 755 stomachs from *M. hubbsi* were examined, of which 475 (62.91%) contained prey items. The TL range of the specimens examined was 14–94 cm (Figure 2). *Merluccius hubbsi* fed almost exclusively on crustaceans in terms of %F, %N and %IRI (Figure 3; Table 1). The euphausiid *Euphausia lucens* was the main prey item by %F, %N and %IRI, but second in terms of %W (Table 1). Among fish, the notothenioid *Patagonotothen ramsayi*, the Argentine anchovy *Engraulis anchoita* and *M. hubbsi* contributed mostly by %W, followed by the cephalopods *Loligo gahi* and



**Fig. 2.** Size–frequency distribution of the Argentine hake, *Merluccius hubbsi*, caught for the diet analysis, by (A) winter inner Patagonian continental shelf (IPCS); (B) winter outer Patagonian continental shelf; (C) summer IPCS. Females (black bars) and males (white bars).



**Fig. 3.** Percentage of index of relative importance (%IRI) for cephalopods, crustaceans and fish in the diet of the Argentine hake *Merluccius hubbsi*, by total length-classes (TL < 34 and TL > 35), during winter on inner Patagonian continental shelf (winter IPCS), winter outer Patagonian continental shelf (winter OPCS) and summer inner Patagonian continental shelf (summer IPCS).

the Argentine squid *Illex argentinus* (Table 1). The TrL of *M. hubbsi* during winter in the OPCS was 3.43.

Of the 1463 specimens examined during summer in the IPCS, 855 (58.44%) contained prey items in their stomachs. The TL of the specimens analysed ranged from 13 cm to 82 cm (Figure 2). During this season, *M. hubbsi* preyed primarily on cephalopods in terms of %F, %W and %IRI (Figure 3), and crustaceans in terms of %N (Table 1). The Argentine squid *Illex argentinus* was the main prey item by %F, %W and %IRI (Table 1). Among crustaceans, the euphausiid *Euphausia lucens*, the hyperiid amphipod *Themisto gaudichaudii* and lobster krill *Munida* spp. were the most important prey (Table 1). The TrL of *M. hubbsi* during summer in the IPCS was 4.51.

### Dietary models


The consumption of *Themisto gaudichaudii*, fish, cephalopods and *Munida* spp. was related to the variables TL, TL<sup>2</sup> and SPTP (GLM 26 in Table 2). On the other hand, the consumption of *Euphausia lucens* and other *M. hubbsi* (by cannibalism) was related to the variables TL and SPTP (GLM 20 in Table 2). The hyperiid amphipods *Themisto gaudichaudii* were more intensively consumed during winter in both OPCS and IPCS, than in summer (Figure 4A; Table 3). The consumption of hyperiid amphipods *T. gaudichaudii* increased from 10 cm to 40 cm predator's TL, and then decreased from 40 cm to 90 cm predator's TL (Figure 4A; Table 3). The euphausiids *Euphausia lucens* were more intensively consumed during winter in OPCS than in IPCS, and during winter in IPCS are also more intensively consumed than in summer (Figure 4B; Table 3). The consumption of the euphausiids *E. lucens* decreased with the predator's length (Figure 4B; Table 3). The lobster krill *Munida* spp. was more intensively consumed during summer in IPCS than during winter in both IPCS and OPCS (Figure 4C; Table 3). The consumption of this species increased from 10 cm to 30 cm and from 30 cm to 60 cm predator's TL in summer IPCS and winter OPCS, respectively, and then decreased as the predator became larger (Figure 4C; Table 3). The fish were more intensively consumed during winter in both OPCS and IPCS than in

summer (Figure 4D; Table 3), and their consumption increased with the predator's TL (Figure 3; Figure 4D; Table 3). Cephalopods were more intensively consumed during summer than in winter in both IPCS and OPCS, and their consumption increased with the predator's length (Figures 3, 4E; Table 3). Cannibalism was higher during winter in IPCS than in OPCS, and it was also higher during winter in OPCS than during summer (Figure 4F; Table 3). The consumption of other *M. hubbsi* increased with the predator's TL (Figure 4F; Table 3).

### DISCUSSION

The Argentine hake *Merluccius hubbsi* of the PS fed mainly upon zooplankton crustaceans, such as euphausiids (*Euphausia lucens*), hyperiid amphipods (*Themisto gaudichaudii*) and lobster krill (*Munida* spp.). The importance of zooplankton crustaceans in the diet of *M. hubbsi* was consistent with earlier observations (Cordo, 1981; Sánchez & Prenski 1996; Ruiz & Fondacaro, 1997; Sánchez & García de la Rosa, 1999; Sánchez, 2009; Ocampo Reinaldo *et al.*, 2011; Temperoni *et al.*, 2013), but differed from others which gave a secondary role to this prey (Angelescu *et al.*, 1958; Angelescu & Cousseau, 1969; Angelescu & Prenski, 1987). High consumption of euphausiids is a common pattern observed in many other gadiform species in the world (Buckley & Livingston, 1997; Garrison & Link, 2000; Orlova *et al.*, 2005), highlighting the importance of this group as a key energetic link between small phytoplankton and large predatory animals. This work also showed that cephalopods (mainly *Illex argentinus*) and fish (*Patagonotothen ramsayi*, *Engraulis anchoita* and other *M. hubbsi*) were also important in some regions and seasons, revealing spatio-temporal changes in the diet of the studied species. These spatio-temporal changes were always accompanied by ontogenetic shifts. On the other hand, the variables sex, maturity stage, age and hour of the day did not influence the consumption of any particular prey item by *M. hubbsi*.

The analysis of the diet of *M. hubbsi* revealed ontogenetic dietary changes associated with size, particularly from



**Fig. 4.** Best models, fitted by generalized linear models for the number of the prey, that explain the changes in the consumption of (A) *Themisto gaudichaudii*; (B) *Euphausia lucens*; (C) *Munida* spp.; (D) fish; (E) cephalopods; (F) *Merluccius hubbsi* (cannibalism). Dotted line, winter IPCS; dashed line winter OPCS; solid line, summer IPCS.

zooplankton (*E. lucens*, *T. gaudichaudii* and *Munida* spp.) to fish and cephalopods. The ontogenetic diet shifts appear to be a universal phenomenon in fish (Wootton, 1990). The diet of fish usually changes because of the morphological changes

that accompany growth (increasing mouth dimensions and stomach capacity, improvement in their locomotion ability, etc.) and allow capture of a broader range of prey sizes and prey types (Wootton, 1990). Large *M. hubbsi* fed on larger,

**Table 3.** Best models that explained the consumption of the most important prey of the Argentine hake *Merluccius hubbsi*. The intercept and coefficient with the standard errors in parentheses (SE) for the variables, the Akaike information criterion (AIC) and the Akaike's weights ( $w$ ) are presented for each model. TL, total length (cm); WinIPCS, winter inner Patagonian continental shelf; WinOPCS, winter outer Patagonian continental shelf.

Prey	Intercept (SE)	Parameters (SE)	AIC	$w$
<i>Themisto gaudichaudii</i>	-1.26 (0.54)	1.37 (0.15)WinIPCS + 1.47 (0.16)WinOPCS + 0.16 (0.03)TL - 0.003 (4×10 <sup>-5</sup> )TL <sup>2</sup>	8135.1	1
<i>Euphausia lucens</i>	2.09 (0.24)	1.75 (0.16)WinIPCS + 3.64 (0.18)WinOPCS - 0.04 (0.005)TL	9360.5	0.72
<i>Munida</i> spp.	-4.05 (1.04)	-1.55 (0.29)WinIPCS - 1.49 (0.33)WinOPCS + 0.22 (0.05)TL - 0.003 (6.7×10 <sup>-4</sup> )TL <sup>2</sup>	1995.2	0.98
Fish	-3.75 (0.39)	0.80 (0.12)WinIPCS + 0.82 (0.12)WinOPCS + 0.08 (0.01)TL - 6.1×10 <sup>-4</sup> (2×10 <sup>-4</sup> )TL <sup>2</sup>	2747	0.98
<i>Merluccius hubbsi</i>	-5.30 (0.32)	1.33 (0.22)WinIPCS + 0.28 (0.27)WinOPCS + 0.05 (0.006)TL	911.3	0.68
Cephalopods	-6.64 (0.48)	-1.65 (0.14)WinIPCS - 1.20 (0.11)WinOPCS + 0.23 (0.02)TL - 0.002 (4×10 <sup>-4</sup> )TL <sup>2</sup>	2454.9	1

more active prey not available to small individuals. In general, *Merluccius* species worldwide feed on zooplankton crustaceans early in their life and with growth, shift their diets to fish and squid (Tanasichuk *et al.*, 1991; Buckley & Livingston, 1997; Garrison & Link, 2000; Carpentieri *et al.*, 2005; Mahe *et al.*, 2007; Cartes *et al.*, 2009).

The feeding on cephalopods and lobster krill *Munida* spp. was higher in summer. These trends are related to seasonal changes of prey abundances, for both cephalopods (Brunetti & Ivanovic, 1992; Brunetti *et al.*, 1998; Crespi *et al.*, 2008) and lobster krill *Munida* spp. (Roux & Piñeiro, 2006), the biomass of which increase during summer on the IPCS. The euphausiid (*Euphausia lucens*) was more intensively consumed during winter on the OPCS, whereas the hyperiid amphipod (*T. gaudichaudii*) was more intensively consumed during winter on the IPCS, in accordance with the distribution of the zooplankton community in the region (Pérez Seijas *et al.*, 1987; Santos, 1994; Sabatini & Alvarez Colombo, 2001; Sabatini, 2008). However, the abundance of euphausiids and amphipods reaches their annual maximum during summer (Pérez Seijas *et al.*, 1987; Sabatini & Alvarez Colombo, 2001) indicating that cephalopods may be preferred to euphausiids and amphipods. Taken together, these patterns suggest a flexible foraging behaviour of *M. hubbsi* which fed on zooplankton throughout the year, but seasonally changed to feed on cephalopods when this kind of prey was available (summer), even when zooplankton was locally abundant. These spatio-temporal variations in accordance with the seasonal fluctuation of prey also supported the idea that *M. hubbsi* has an opportunistic or adaptive foraging behaviour. This kind of adaptive foraging behaviour may make a species relatively resilient to fisheries-induced ecosystem changes (Dunn *et al.*, 2013).

The quality of food is essential to understanding the energy sources consumed by predators (Vollenweider *et al.*, 2011), and should be considered when spatio-temporal and ontogenetic shifts of *M. hubbsi* are discussed. Cephalopods were preferred by hake when this prey is accessible. Moreover, cephalopods and fish were important prey by weight, whereas zooplankton crustaceans were important by number. The high energy provided by cephalopods and fish in relation to zooplankton crustaceans (Ciancio *et al.*, 2007) justifies their choice. Based on the proximate composition, fish and cephalopods are considered high quality prey (4–7 J g<sup>-1</sup>), while zooplankton crustaceans such as euphausiids and amphipods are low quality prey (2–3 J g<sup>-1</sup>) (Ciancio *et al.*, 2007). The high proportion of chitin from the exoskeletons of zooplankton crustaceans also determines preference for cephalopods and fish by larger hake. However, prey selection also depends of prey mobility, prey size and easy access

(Wootton, 1990) conferring higher profitability of zooplankton crustaceans to the small *M. hubbsi*.

A controversial component in the diet of *M. hubbsi* was other hake. Cannibalism increased with predator's length, and contributed a maximum of 8.14% to the diet during winter on the IPCS. Gadiforms, particularly hakes, are known to be highly cannibalistic (Juanes, 2003). Chiou *et al.* (2006) found that cannibalism would increase when the population of cannibals is abundant, or when other food sources are insufficient. Previous authors reported higher rates of cannibalism in the diet of *M. hubbsi* than that observed in the present work (Cordo, 1981; Sánchez & García de la Rosa, 1999; Ocampo Reinaldo *et al.*, 2011), whereas others suggested that cannibalism occurs when food is scarce and as a secondary prey (Angelescu & Prenske, 1987; Sánchez & Prenske, 1996; Ruiz & Fondacaro, 1997; Sánchez, 2009). During winter on the IPCS the lowest biomasses of the most important prey of *M. hubbsi* occur (Pérez Seijas *et al.*, 1987; Brunetti *et al.*, 1998; Sabatini & Alvarez Colombo, 2001; Roux & Piñeiro, 2006; Crespi *et al.*, 2008), and there is a high overlap of different age and size-classes (Macchi *et al.*, 2007). The strong cannibalism during winter on the IPCS suggests that it is related to a combination of factors, such as the abundance and availability of alternative prey and to the overlap of different size-classes.

This paper provides the first estimation of the trophic level of *M. hubbsi*, which occupies different trophic positions according to season and region. During winter *M. hubbsi* can be considered a secondary consumer (3 < TrL < 4) due to the high consumption of zooplankton crustaceans. On the other hand, it can be considered a tertiary consumer, or apex predator (TrL > 4), during summer when the cephalopod *I. argentinus* is the main prey. The few works of worldwide extent that have estimated the TrL of *Merluccius* spp. also have shown that it ranged from 3.2 to 4.05 (Stergiou & Karpouzi, 2002; Iitembu *et al.*, 2012). The spatio-temporal variations observed in TrL may be a result of the variability of the species dominance on the study area (Pérez Seijas *et al.*, 1987; Brunetti *et al.*, 1998; Sabatini & Alvarez Colombo, 2001; Roux & Piñeiro, 2006; Crespi *et al.*, 2008). Since exploited marine ecosystems are characterized by a decline of the mean TrL in landings (Pauly *et al.*, 1998), the monitoring of the TrL, the food web structure and fisheries landings are needed on the PS for appropriate management in future years.

In contrast to other gadoids, such as cod, which have been harvested for centuries, large scale hake fishing is relatively recent (Pitcher & Alheit, 1995). However, in Argentina the estimated total biomass in northern stock declined from about 690,000 tons in 1986 to 287,500 tons in 1999, whereas the PS was reduced from 1,520,000 tons in 1993 to

844,500 tons in 1999 (Aubone *et al.*, 2004). A permanent closed zone covering a great part of the Argentine Sea has been established since 1998 to protect *M. hubbsi* of the PS. In this context, fishery researchers need basic diet composition data for each predator to develop ecosystem models involving energy (Cochrane, 2002; Thrush & Dayton, 2010). From a viewpoint of population dynamics, distinguishing among the different causes of mortality including cannibalism (and if it is uniform across the study area) is essential to accurately reflect the real situation and to make fisheries management more effective (Hollowed *et al.*, 2000). Our results provide the diet data requirements that seem to be one among many necessary steps for developing ecosystem models.

## ACKNOWLEDGEMENTS

We thank the Instituto Nacional de Investigación y Desarrollo Pesquero (INIDEP) and the scientists on-board the research vessels (Claudia Dato, Juampi Simonazzi, Hugo Brachetta, Claudio Buratti and Luis Peyrot) for the stomachs collected. The librarians from INIDEP (Gaby Cosulich, Gaby Silvoni, Leticia Lizondo and Quimey Navas) who provided the most helpful documentation. We are also grateful to Brenda Temperoni and Enrique Boschi for their help in identifying crustaceans.

## FINANCIAL SUPPORT

Belleggia Mauro was supported by a post-doctoral scholarship from CONICET. This paper is an INIDEP contribution number 1860.

## REFERENCES

- Acha E.M., Mianzan H.W., Guerrero R.A., Favero M. and Bava J. (2004) Marine fronts at the continental shelves of austral South America. Physical and ecological processes. *Journal of Marine Systems* 44, 83–105.
- Angelescu V. and Cousseau M.B. (1969) Alimentación de la merluza en la región del talud continental Argentino, época invernal (*Merluccidae Merluccius hubbsi*). *Boletín del Instituto de Biología Marina* 19, 5–84.
- Angelescu V., Gneri F.S. and Nani A. (1958) *La merluza del mar Argentino (biología y taxonomía)*. Buenos Aires: Secretaría Marina, Servicio de Hidrografía Naval.
- Angelescu V. and Prenske L.B. (1987) Ecología trófica de la merluza común del mar Argentino (*Merluccidae, Merluccius hubbsi*). Parte 2. Dinámica de la alimentación analizada sobre la base de las condiciones ambientales, la estructura y las evaluaciones de los efectivos en su área de distribución. *Contribución INIDEP* 561, 1–205.
- Arkhipkin A.I., Middleton D.A.J., Portela J.M. and Bellido J.M. (2003) Alternative usage of common feeding grounds by large predators: the case of two hakes (*Merluccius hubbsi* and *M. australis*) in the Southwest Atlantic. *Aquatic Living Resources* 16, 487–500.
- Aubone A., Bezzi S.I., Cañete G., Castrucci R., Dato C., Irusta G., Madirolas A., Pérez M., Renzi M., Santos B., Simonazzi M. and Villarino M.F. (2004) Evaluación y sugerencias de manejo del recurso merluza (*Merluccius hubbsi*). La situación hasta 1999. In Sánchez R. and Bezzi S. (eds) *El Mar Argentino y sus recursos pesqueros. Tomo IV. Los peces marinos de interés pesquero. Caracterización biológica y evaluación del estado del estado de explotación*. Mar del Plata: INIDEP, pp. 207–235.
- Bastida R.O. and Torti M.R. (1973) Los Isópodos Serolidae en la Argentina. Clave para su reconocimiento. *Physis* 84, 19–46.
- Bastida R.O., Roux A. and Martínez D.E. (1992) Benthic communities of the Argentine continental shelf. *Oceanologica Acta* 15, 687–698.
- Bezzi S.I., Verazay G.A. and Dato C.V. (1995) Biology and fisheries of Argentine hakes (*M. hubbsi* and *M. australis*). In Alheit J. and Pitcher T.J. (eds) *Hake: biology, fisheries and markets*. London: Chapman & Hall, pp. 229–267.
- Brunetti N.E. and Ivanovic M.L. (1992) Distribution and abundance of early life stages of squid (*Illex argentinus*) in the south-west Atlantic. *ICES Journal of Marine Science* 49, 175–183.
- Brunetti N.E., Ivanovic M., Rossi G., Elena B. and Pineda S. (1998) Fishery biology and life history of *Illex argentinus*. In Okutani T. (ed.) *Contributed papers to International Symposium on Large Pelagic Squid*. Tokyo: Marine Fishery Resources Research Center, pp. 217–232.
- Buckley T.W. and Livingston P.A. (1997) Geographic variation in the diet of Pacific hake with a note on cannibalism. *California Cooperative Oceanic Fisheries Investigations* 38, 53–62.
- Carpentieri P., Colloca F., Cardinale M., Belluscio A. and Ardizzone G.D. (2005) Feeding habits of European hake (*Merluccius merluccius*) in the central Mediterranean Sea. *Fishery Bulletin* 103, 411–416.
- Cartes J.E., Hidalgo M., Papiol V., Massutí E. and Moranta J. (2009) Changes in the diet and feeding of the hake *Merluccius merluccius* at the shelf-break of the Balearic Islands: influence of the mesopelagic-boundary community. *Deep-Sea Research I* 56, 344–365.
- Chiu W., Chen C., Wang C. and Chen C. (2006) Food and feeding habits of ribbonfish *Trichiurus lepturus* in coastal waters of south-western Taiwan. *Fisheries Science* 72, 373–381.
- Ciancio J., Pascual M. and Beauchamp D. (2007) Energy density of Patagonian aquatic organisms and empirical predictions based on water content. *Transactions of the American Fisheries Society* 136, 1415–1422.
- Cochrane K.L. (2002) The use of scientific information in the design of management strategies; Chapter 5. Rome: FAO, Fisheries Technical Paper No. 424, pp. 95–130.
- Cordo H.D. (1981) Resultados sobre la alimentación de la merluza del mar epicontinental Argentino (*Merluccius hubbsi*). Análisis biológico y estadístico de los datos obtenidos de las campañas de B/I “Shinkai Maru” y “Walther Herwig” (1978–1979). In Angelescu V. (ed.) *Campañas de investigación pesquera realizadas en el mar Argentino por los B/I “Shinkai Maru” y “Walther Herwig”, años 1978 y 1979. Resultados de la parte argentina*. Mar del Plata: INIDEP, pp. 299–312.
- Cortés E. (1997) A critical review of methods of studying fish feeding based on analysis of stomach contents: application to elasmobranch fishes. *Canadian Journal of Fisheries and Aquatic Sciences* 54, 726–738.
- Cortés E. (1999) Standardized diet compositions and trophic levels of sharks. *ICES Journal of Marine Science* 56, 707–717.
- Cousseau M.B. and Perrota R.G. (2004) *Peces Marinos de Argentina: Biología, Distribución y Pesca*. Mar del Plata: INIDEP.
- Crawley M.J. (2005) *Statistics: an introduction using R*. Chichester: John Wiley & Sons.
- Crespi A.C.A., Morsan E.M. and Barón P.J. (2008) Contribution to understanding the population structure and maturation of *Illex argentinus* (Castellanos, 1960): the case of the inner-shelf spawning groups in San Matias Gulf (Patagonia, Argentina). *Journal of Shellfish Research* 27, 1225–1231.



- Di Giacomo E.E., Calvo J., Perier M.R. and Morriconi E.R. (1993) Spawning aggregations of *Merluccius hubbsi*, in Patagonian waters: evidence for a single stock? *Fisheries Research* 16, 9–16.
- Dunn M.R., Stevens D.S., Forman J.S. and Connell A. (2013) Trophic interactions and distribution of some Squaliforme sharks, including new diet descriptions for *Deania calcea* and *Squalus acanthias*. *PLoS ONE* 8, e59938. doi:10.1371/journal.pone.0059938.
- Ebert D.A. and Bizzarro J.J. (2007) Standardized diet compositions and trophic levels of skates (Chondrichthyes: Rajiformes: Rajoidei). *Environmental Biology of Fishes* 80, 221–237.
- Ehrlich M.D. and Ciechomski J.D. (1994) Reseña sobre la distribución de huevos y larvas de merluza (*Merluccius hubbsi*) basada en veinte años de investigación. *Frente Marítimo* 15, 37–50.
- Franklin A.B., Shenk T.M., Anderson D.R. and Burnham K.P. (2001) Statistical model selection: the alternative to null hypothesis testing. In Shenk T.M. and Franklin A.B. (eds) *Modeling in natural resource management: development, interpretation, and application*. Washington, DC: Island Press, pp. 75–90.
- Garrison L.P. and Link J. (2000) Diets of five hake species in the northeast United States continental shelf ecosystem. *Marine Ecology Progress Series* 204, 243–255.
- Guerrero R.A. and Piola R.A. (1997) Masas de agua de la plataforma continental. In Boschi E.E. (ed.) *El Mar Argentino y sus Recursos Pesqueros*. Volume I. Mar del Plata: INIDEP, pp. 107–118.
- Hollowed A.B., Ianelli J.N. and Livingston P.A. (2000) Including predation mortality in stock assessments: a case study for Gulf of Alaska walleye Pollock. *ICES Journal of Marine Science* 57, 279–293.
- Itembu J.A., Miller T.W., Ohmori K., Kanime A. and Wells S. (2012) Comparison of ontogenetic trophic shift in two hake species, *Merluccius capensis* and *Merluccius paradoxus*, from the Northern Benguela Current ecosystem (Namibia) using stable isotope analysis. *Fisheries Oceanography* 21, 215–225.
- Johnson J.B. and Omland K.S. (2004) Model selection in ecology and evolution. *Trends in Ecology and Evolution* 19, 101–108.
- Juanes F. (2003) The allometry of cannibalism in piscivorous fishes. *Canadian Journal of Fisheries and Aquatic Sciences* 60, 594–602.
- Laptikhovsky V. and Fetisov A. (1999) Scavenging by fish of discards from the Patagonian squid fishery. *Fisheries Research* 41, 93–97.
- Lloris D., Matallanas J. and Oliver P. (2003) *Merluzas del Mundo (Familia Merlucciidae)*. Catálogo comentado e ilustrado de las merluzas conocidas. Rome: FAO.
- Lucifora L.O., García V.B. and Escalante A.H. (2009) How can the feeding habits of the sand tiger shark influence the success of conservation programs? *Animal Conservation* 12, 291–301.
- Macchi G.J. and Pájaro M. (2003) Comparative reproductive biology of some commercial marine fishes from Argentina. *Fisken og Havet* 12, 69–77.
- Macchi G.J., Pájaro M. and Dato C. (2007) Spatial variations of the Argentine hake (*Merluccius hubbsi* (Marini, 1933)) spawning shoals in the Patagonian area during a reproductive season. *Revista de Biología Marina y Oceanografía* 42, 345–356.
- Mahe K., Amara R., Brychaert T., Kacher M. and Brylinski J.M. (2007) Ontogenetic and spatial variation in the diet of hake (*Merluccius merluccius*) in the Bay of Biscay and the Celtic Sea. *ICES Journal of Marine Science* 64, 1210–1219.
- Nelson J.S. (2006) *Fishes of the world*. New York: John Wiley and Sons.
- Ocampo Reinaldo M., González R. and Romero M.A. (2011) Feeding strategy and cannibalism of the Argentine hake *Merluccius hubbsi*. *Journal of Fish Biology* 79, 1795–1814.
- Orlova E.L., Dolgov A.V., Rudneva G.B. and Nesterova V.N. (2005) The effect of abiotic and biotic factors on the importance of macroplankton in the diet of Northeast Arctic cod in recent years. *ICES Journal of Marine Science* 62, 1463–1474.
- Pauly D., Christensen V., Dalsgaard J., Froese R. and Torres F. Jr (1998) Fishing down marine food webs. *Science* 279, 860–863.
- Pérez Seijas G.M., Ramírez F.C. and Viñas M.D. (1987) Variaciones de la abundancia numérica y biomasa del zooplancton de red en el Golfo San Jorge (Año 1985). *Revista de Investigación y Desarrollo Pesquero* 7, 5–20.
- Pinkas L.M., Oliphant S. and Iverson I.L.K. (1971) Food habits of albacore, bluefin tuna and bonito in Californian waters. *California Fish and Game* 152, 1–105.
- Piola A.R., Avellaneda N.M., Guerrero R.A., Jardón F.P., Palma E.D. and Romero S.I. (2010) Malvinas-slope water intrusions on the northern Patagonia continental shelf. *Ocean Science Discussions* 6, 2939–2974.
- Pitcher T.J. and Alheit J. (1995) What makes a hake? A review of the critical biological features that sustain global hake fisheries. In Alheit J. and Pitcher T.J. (eds) *Hake: biology, fisheries and markets*. London: Chapman & Hall, pp. 1–14.
- Ramírez F.C. and Viñas M.D. (1985) Hyperiid amphipods found in Argentine shelf waters. *Physis* 43, 25–37.
- Ramírez F.C. (1971) Eufáusidos de algunos sectores del Atlántico Sud-Occidental. *Physis* 30, 385–405.
- Renzi M.A. and Pérez M.A. (1992) Un criterio para la determinación de la edad en juveniles de merluza (*Merluccius hubbsi*) mediante la lectura de otolitos. *Frente Marítimo* 11, 15–31.
- Roux A. and Piñero R. (2006) Crustáceos decápodos y estomatópodos asociados a la pesquería del langostino patagónico *Pleoticus muelleri* (Bate, 1888) en el Golfo San Jorge, Argentina. *Revista de Investigación y Desarrollo Pesquero* 18, 33–43.
- Ruiz A.E. and Fondacaro R.R. (1997) Diet of hake (*Merluccius hubbsi* Marini) in a spawning and nursery area within Patagonian shelf waters. *Fisheries Research* 30, 157–160.
- Sabatini M.E. (2008) El ecosistema de la plataforma patagónica austral, marzo-abril 2000. Composición, abundancia y distribución del zooplancton. *Revista de Investigación y Desarrollo Pesquero* 19, 5–21.
- Sabatini M.E. and Álvarez Colombo G.L. (2001) Seasonal pattern of zooplankton biomass in the Argentinian shelf off Southern Patagonia (45°–55°S). *Scientia Marina* 65, 21–31.
- Sánchez F. (2009) Alimentación de la merluza (*Merluccius hubbsi*) en el Golfo San Jorge y aguas adyacentes. *Informe Técnico INIDEP* 75, 1–21.
- Sánchez F. and Prenski L.B. (1996) Ecología trófica de peces demersales en el Golfo San Jorge. *Revista de Investigación y Desarrollo Pesquero* 10, 57–71.
- Sánchez F. and García de la Rosa S.B. (1999) Alimentación de *Merluccius hubbsi* e impacto del canibalismo en la región comprendida entre 34°50'S–47°S del Atlántico Sudoccidental. *Revista de Investigación y Desarrollo Pesquero* 12, 77–93.
- Santos B.A. (1994) Biomasa y composición del zooplancton de interés trófico-pesquero en áreas del Atlántico Sudoccidental. Campaña H-01/93, 42°–51°S, enero 1993. *Informe Técnico INIDEP* 2, 5–21.
- Sea Around Us (2014) *A global database on marine fisheries and ecosystems*. <http://www.seaaroundus.org> (accessed 15 April 2014).
- Stergiou K.I. and Karpouzi V.S. (2002) Feeding habits and trophic levels of Mediterranean fish. *Reviews in Fish Biology and Fisheries* 11, 217–254.

- Tanasichuk K.W., Ware D.M., Shaw W. and McFarlane G.A.** (1991) Variation in diet, daily ration, and feeding periodicity of Pacific hake (*Merluccius productus*) and spiny dogfish (*Squalus acanthias*) off the lower west coast of Vancouver Island. *Canadian Journal of Fisheries and Aquatic Sciences* 48, 2118–2128.
- Temperoni B., Viñas M.D. and Buratti C.C.** (2013) Feeding strategy of juvenile (age-0+ year) Argentine hake *Merluccius hubbsi* in the Patagonian nursery ground. *Journal of Fish Biology* 83, 1354–1370.
- Thrush S.F. and Dayton P.K.** (2010) What can ecology contribute to ecosystem-based management? *Annual Review of Marine Science* 2, 419–441.
- Vögler R., Milessi A.C. and Duarte L.O.** (2009) Changes in trophic level of *Squatina guggenheim* with increasing body length: relationships with type, size and trophic level of its prey. *Environmental Biology of Fishes* 84, 41–52.
- Vollenweider J., Heintz R., Schaufler L. and Bradshaw R.** (2011) Seasonal cycles in whole-body proximate composition and energy content of forage fish vary with water depth. *Marine Biology* 158, 413–427.
- and
- Wootton R.J.** (1990) *Ecology of teleost fishes*. London, UK: Chapman & Hall
- Correspondence should be addressed to:**  
M. Belleggia  
Instituto Nacional de Investigación y Desarrollo Pesquero  
Paseo Victoria Ocampo 1, Mar del Plata, B7602HSA  
Argentina  
email: [belleggia@inidep.edu.ar](mailto:belleggia@inidep.edu.ar)